



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁴ : C07K 3/00, 13/00, A61K 9/50 A61K 35/48 // A61K 37/04 C12N 9/36, 9/96	A2	(11) International Publication Number: WO 87/ 05300 (43) International Publication Date: 11 September 1987 (11.09.87)
(21) International Application Number: PCT/GB87/00143 (22) International Filing Date: 27 February 1987 (27.02.87) (31) Priority Application Number: 8604983 (32) Priority Date: 28 February 1986 (28.02.86) (33) Priority Country: GB (71) Applicant (for all designated States except US): BIOCOMPATIBLES LIMITED [GB/GB]; 2 Gray's Inn Square, London WC1R 5AF (GB). (72) Inventors; and (75) Inventors/Applicants (for US only) : HAYWARD, James, Arthur [US/US]; 4 Chuck Court, Port Jefferson, NY 11777 (US). JOHNSTON, David, Samuel [GB/GB]; 65 Fitzjohns Avenue, Hampstead, London NW3 6PE (GB).		(74) Agents: CRESSWELL, Thomas, Anthony et al.; J.A. Kemp & Co., 14 South Square, Gray's Inn, London WC1R 5EU (GB). (81) Designated States: AT (European patent), BE (European patent), CH (European patent), DE (European patent), FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent), US. Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: PROTEIN PRESERVATION (57) Abstract A process for preserving a material having water-dependant structure comprising contacting the material with an aqueous solution of a polyhydroxy compound then removing water from the material.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

-1-

PROTEIN PRESERVATION

The present invention relates to a method of dry storage which now permits the long-term preservation of materials such as haemoglobin, erythrocytes, liposomes and cells.

Oxidation of the iron atom of the haem prosthetic group from 2+ to the 3+ state converts oxy- or deoxyhaemoglobin to the biologically nonfunctional methaemoglobin. The reaction is promoted by conditions of low oxygen tension and is minimised in vivo by the presence of potent reducing systems operating in the cytoplasm of intact erythrocytes. When erythrocytes are obtained from human donors for transfusion, their storage ex vivo is limited, in part, by the accelerated formation of oxidised haem (methaemoglobin).

Storage in the dry state limits hydrolysis and prevents some of the oxidative damage that can destroy chemical or biological activity. When dealing with biological macromolecules, however, water removal can irreversibly alter secondary, tertiary or quaternary structures and thereby eliminate biological efficacies.

Dehydration of biological membranes is usually accompanied by irrevocable loss of structural and functional integrity (Crowe, Crowe and Jackson, 1983). Upon rehydration of biological membranes morphological

-2-

damage is apparent in the redistribution of intramembraneous particles and the fusion between membranes. Moreover, in those membrane systems that are associated with a specific enzyme activity (such as the transport of Ca^{++}), rehydration induces a complete loss of activity.

Artificial membrane systems prepared by dispersion of phospholipids in aqueous solutions, have received much attention because of their utility in basic research and their potential applications in biotechnology. Dispersions of phospholipids spontaneously form sealed structures called liposomes - the encapsulated volume within a liposome enables the liposome to function as a pharmacological "capsule" or artificial cell. Among the solutes that have been encapsulated in liposomes is human haemoglobin. The "haemosomes" so generated hold much utilitarian promise as erythrocyte surrogates (Djordjevic and Miller, 1980; Gaber et al., 1983; Hayward et al., 1985). However when liposomes containing entrapped solutes are subjected to dehydration/rehydration, the solutes are lost to the bulk solution (Madden et al., 1985).

The effect of encapsulation of haemoglobin within a phospholipid liposome was to catalyze the oxidation of haemoglobin upon rehydration. No oxyhaemoglobin could be regenerated from rehydrated haemosomes. Moreover, much of the visible absorbance of the rehydrated samples was bleached, resulting in essentially colourless samples.

SUBSTITUTE SHEET

-3-

These observations are similar to those described by Szebeni et al. (1985), who found that unsaturated phospholipids promoted haemoglobin oxidation and were capable of extracting the haem away from the globin portion of the molecule.

We have found that by drying proteins from aqueous solutions containing saccharides, chemical and physical degradative processes can be avoided. If the carbohydrate selected is itself a solid, then a compact, solid protein-carbohydrate mixture is obtained which can be repeatedly cooled below 0°C and warmed to room temperature without diminishing enzyme activity. Moreover this technique may be extended to facilitate storage of other materials having water-dependant structure such as liposomes and even intact cells.

We have examined the ability of trehalose and other sugars to preserve after dehydration/rehydration the biological activity of isolated proteins, protein systems, biological and artificial membranes, and whole cells.

The present invention therefore provides a process for preserving a material having a water-dependant structure comprising contacting the material with an aqueous solution of a polyhydroxy compound then removing water from the material.

By "material having a water-dependant structure" it is intended to encompass such materials as proteins, liposomes and cells which have secondary, tertiary or

SUBSTITUTE SHEET

-4-

quaternary structure determined by hydrophobic and/or hydrophilic interactions in the presence of water which structure is irreversibly altered or partially or completely destroyed by removal of the water.

By "polyhydroxy compound" it is intended to encompass any material having two or more hydroxy groups per molecule which is capable of hydrogen bonding to, or otherwise stabilising, a material having a water-dependant structure. Suitable polyhydroxy compounds include sugars and similar or related polyols, especially trioses, pentoses and hexoses, mono-, di or oligosaccharides and related polyols. Preferred polyhydroxy compounds are glucose, galactose and non-reducing sugars especially trehalose.

The present invention, in a particular embodiment also provides a process for preserving a material having a water-dependant structure which encapsulates one or more spaces containing aqueous solutions or suspensions which comprises contacting the material with, and introducing into one or more encapsulated spaces therein, an aqueous solution of polyhydroxy compound then removing water from the material.

The present invention in another aspect provides a dry formulation comprising a material having a water-dependant structure and a polyhydroxy compound. The polyhydroxy compound will be in intimate admixture with the material having a water-dependant structure and, in the

SUBSTITUTE SHEET

-5-

case where the material encapsulates one or more spaces containing the residue of aqueous solutions or suspensions, preferably the polyhydroxy compound is also dispersed within those spaces.

The preservation process of the present invention is particularly suitable for the storage of haemoglobin with minimal oxidation of the heme-iron and with retention of gas-transporting capacity upon rehydration. The dry haemoglobin may be prepared from simple aqueous solutions of purified haemoglobin, from erythrocyte lysates, from intact erythrocytes or from erythrocyte surrogates (haemosomes).

Whilst not wishing to be bound by any theory it is postulated that the stabilisation afforded according to the present invention may be related to that observed in certain instances of "anhydrobiosis".

The organisation of many biological structures depends absolutely upon the presence of water. Hydrophobic structures, such as the membranes which delimit all cells and their organelles, arise by the expulsion of water from their non-polar parts. Hydrophilic structures such as globular proteins are stabilised by hydrogen-bonding with water in solution and by the maintenance of specific dipoles in a medium of specific dielectric constant.

Some organisms, however, possess a remarkable capacity to survive near-total dehydration, and they may exist in the dehydrated states for periods measured in

SUBSTITUTE SHEET

-6-

years. This fact was recognised as long ago as the year 1800 when it was found that in a dried state tardigrades, small sand-living organisms, were able to withstand temperatures well below freezing and as hot as boiling water (Crowe and Clegg, 1973 and 1978). Normal metabolic processes resumed in these organisms within thirty minutes of their rehydration even after decades of dry storage.

In recent years the ability to survive desiccation, now termed "anhydrobiosis", has been shown to be a property of a small but diverse group of organisms including fungal spores, macrocysts of the slime mold Dictyostelium, baker's yeast, brine shrimp cysts, nematodes and plant seeds (for review see Crowe and Crowe, 1984).

Trehalose, a non-reducing disaccharide of glucose that is distributed widely in nature, has been found at particularly high concentrations (as much as 20% of the dry weight) in several of the organisms capable of anhydrobiosis. The survival of dehydration by some organisms is correlated with the synthesis of trehalose during dehydration (Madin and Crowe, 1975) or its degradation following rehydration (Clegg, 1964). This molecule was implicated in the "water replacement hypothesis", which suggested that the hydroxyl groups of simple carbohydrates might replace the hydration shell of the polar head groups of phospholipids that is lost during dehydration (Crowe and Clegg, 1973).

Recent experiments using infrared spectroscopy

SUBSTITUTE SHEET

-7-

(Crowe, Crowe and Chapman, 1984), have provided evidence that trehalose may substitute for water by forming hydrogen bonds with the head groups of phospholipid bilayers thereby stabilising membrane structure in the absence of water. Support for this model was obtained by Madden et al. (1985) who demonstrated that the integrity of unilamellar liposomes is retained (i.e. no leakage of solute from the internal aqueous compartment of the liposome) if dehydration is carried out in the presence of carbohydrate. This protective capacity was shared among all of the carbohydrates tested.

Sugars and similar poly alcohols have been utilised in solution to preserve enzymatic activities (Bradbury and Jakoby, 1972), inhibit irreversible aggregation (Frigon and Lee, 1972) and increase the temperature at which proteins are thermally denatured (Gerlisma, 1968; Neucere and St. Angelo, 1972).

Due to the widespread use of proteins in analytical, synthetic and biological chemistry, we envisage that carbohydrate-mediated protection of dried proteins will have many applications. The storage stability of most proteins should be enhanced by the processes we describe.

A particularly novel extension of our invention concerns the stabilisation of dehydrated cells and their surrogates. It is possible to transiently produce pores in both biological and artificial membrane which spontaneously reseal. During the time course of their existence,

SUBSTITUTE SHEET

-8-

however, these pores are of sufficient diameter to permit the entry of carbohydrates into the cytoplasm (cells) or encapsulated volume (liposomes). Thus, it should be possible to introduce trehalose (or other carbohydrates) into the cytoplasm of human erythrocytes, enabling their extended storage in the dehydrated state. Autologous red cells could therefore be stored indefinitely for patients at high risk of anaemia. Similarly, haemosomes could be prepared and stored in bulk following dehydration. Application of dry haemosomes would be a simple matter following their rehydration ad hoc.

The invention will now be illustrated by non-limiting Examples and with reference to the accompanying drawings in which; Fig. 1 is a histogram showing Percentages of methaemoglobin formed upon rehydration of haemoglobin powders. Sugars were present at concentrations of 0.25 molar following rehydration; haemoglobin was 0.015 molar. 1) Oxyhaemoglobin without sugar, 2) Deoxyhaemoglobin without sugar, and 3) to 11) Oxyhaemoglobin with 3) Arabinose, 4) Glactose, 5) Fucose, 6) Glucose, 7) Mannose, 8) Maltose, 9) lactose, 10) Trehalose and 11) Sucrose.

EXAMPLES 1 AND 2

Different sugars are utilised to produce dry powders of two separate proteins, chicken egg-white

SUBSTITUTE SHEET

-9-

lysozyme (Example 1) and human haemoglobin (Example 2). Structural stability was examined by calorimetry of the wet and dry lysozyme preparations. Functional stability was verified by spectroscopic evaluation of gas transport by haemoglobin.

Carbohydrate-protein mixtures were prepared in solution by addition of carbohydrate to the protein dissolved in water or buffer. A variety of trioses, pentoses and hexoses were examined as well as mono- and disaccharides. Concentrations of the sugars were varied, from 1:1 (weight:weight) up to a maximum of 0.25 molar sugar.

Water was removed by one of three methods: a) evaporation, in which the solution is mildly heated under a steady stream of gaseous nitrogen; b) evacuation, in which the sample is exposed to reduced pressure at room temperature, or, c) lyophilisation, in which the solution is first frozen in liquid nitrogen and the water sublimed under high vacuum. Dried samples were stored for extended period over phosphorus pentoxide in vacuo.

Calorimetric evaluation was performed by differential scanning calorimetry (DSC) using Perkin-Elmer DSC-2. The heating rate was 2.5°C/minute and a full-scale deflection on the recorder corresponded to a change in heat flow of 0.25m.cal./sec. The transition temperatures obtained were the temperatures of maximal heat flow.

Table 1 contains the denaturation temperatures

SUBSTITUTE SHEET

-10-

and denaturation enthalpy and entropy changes of lysozyme dissolved in water and in solutions of either of the sugars glucose, galactose or trehalose. The presence of any of the three sugars increases the denaturation temperature, in general agreement with the observations reported by others (Gerlsma, 1968; Neucere and St. Angelo, 1972). Hence, in solution, the sugars stabilise the native structure of the protein with respect to the denatured state. This property is exhibited by each of the three sugars examined.

SUBSTITUTE SHEET

-11-

TABLE ONE

DENATURATION TEMPERATURES AND DENATURATION ENTHALPY AND
ENTROPY CHANGES OF LYSOZYME IN PURE WATER AND AQUEOUS

SUGAR SOLUTIONS

	Concn. Lysozyme in Water (w/wt)	Sugar	Ratio Sugar to Lysozyme (mole:mole)	$T_d(^{\circ}\text{C})$	ΔS_{cal} (kcal.-mole) (cals.-kmole)	
					ΔH_{cal}	ΔS_{cal}
a	5	None	-	72.8	142	410.5
b	10	"	"	70.9	116.2	337.7
c	20	"	"	68.8	105.7	309.1
d	10	Glucose	1:1	73.8	118.9	342.7
e	"	Trehalose	"	73.6	119.6	344.9
f	"	Galactose	"	73.5	139.0	401.0
g	"	"	"	73.6	133.0	384.2
h	5	"	"	73.9	136.3	392.7
i	20	"	"	75.3	101.0	289.6
j	10	Glucose	5:1	83.2	88.8	249.8
k	"	Trehalose	"	79.8	111.5	315.8
l	"	Galactose	"	81.5	90.4	254.9

SUBSTITUTE SHEET

-12-

The temperatures of maximum heat flow and the enthalpy and entropy changes associated with the transitions for freeze-dried lysozyme in the presence or absence of carbohydrate are outlined in Table 2. The endothermic transition of pure dry lysozyme occurs at a temperature approximately 70°C above the denaturation temperature of the protein in water. In the solid state the polypeptide chains of a protein will be less flexible than in solution at the same temperature.

Therefore, a dry protein will reach the degree of flexibility necessary for unfolding of the protein structure at a higher temperature than a protein in aqueous solution.

SUBSTITUTE SHEET

-13-

TABLE TWO

TRANSITION TEMPERATURES AND TRANSITION ENTHALPY AND ENTROPY
CHANGES OF FREEZE DRIED LYSOZYME AND SUGAR-LYSOZYME MIXTURES

Ratio Sugar to Lysozyme (mole:mole)	Sugar	Mean T_{tr} ($^{\circ}$ C)	Mean ΔH_{cal} (kcal.-mole)	S.D./ Range	S.D./ Range	ΔS_{cal} (cals.-kmole)	number of tests
a	1:1 Glucose	117.0	152.6	3.56	13.1	391.1	8
b	" Trehalose	127.3	123.8	5.56	7.6	309.2	11
c	" Galactose	128.1*	368.0	3.25	22.1	917.2	6
		110.9	66.0	3.45	4.1	171.9	
d	5:1 Glucose	122.5	348.3	123.7- 120.9	351.7- 344.9	880.3	3
e	" Trehalose	139.4	140.1	1.65	13.9	339.6	4
f	Pure lysozyme	139.6	63.9	5.80	13.9	154.8	9
g	Pure lysozyme (freeze dried)	128.8	49.8	9.80	11.1	123.9	8
x	Thermogram displays two separate endotherms						

SUBSTITUTE SHEET

-14-

In the presence of sugar, the transition temperature is decreased, and the enthalpy and entropy changes for the transition equal or exceed those values seen for the protein in solution. Thus, it is apparent that as the water of hydration is removed from the surface of the protein, it is gradually replaced by sugar, resulting in a stabilisation of protein conformation that does not occur in the absence of added sugar.

Although each of the three sugars studied was capable of restoring the calorimetric behaviour of dissolved lysozyme to the dried protein, it was observed that the heated powders containing glucose or galactose were "browned". This browning reaction did not occur in the presence of trehalose. Thus, the selective pressure toward the occurrence of trehalose in anhydrobiotic organisms may be due less to its capacity to substitute for water than to its non-reducing characteristics which prevent the browning reaction. Nonetheless, in our invention any of the above-mentioned sugars are appropriate for storage of dry protein-sugar mixtures.

These results were extended by determinations of the functional integrity of a human haemoglobin before and after dehydration/rehydration in the presence or absence of various carbohydrates. The relative concentrations of oxy-, deoxy- and methaemoglobins were calculated from the spectra obtained in solution according to the methods described in Benesch et al. (1965).

SUBSTITUTE SHEET

-15-

In Figure 1 are presented the percentages of methaemoglobin formed in solutions of human haemoglobin following a cycle of dehydration/rehydration. When dried haemoglobin without sugar was rehydrated, the solution assumed the brown colour characteristic of methaemoglobin. The percentage of methaemoglobin formed increased with the duration of storage, up to 90% of the total haem. Similar spectra could be generated by catalytic oxidation of the haem after addition of potassium ferricyanide.

In strong contrast to the behaviour of rehydrated haemoglobin in the absence of sugar, each of the carbohydrates tested was capable of protecting haemoglobin against oxidative damage. The protective effect was clearly manifest even in those samples that had been stored in the dried state for periods in excess of three months. Rehydrated sugar-haemoglobin mixtures were capable of reversible gas exchange as evidenced by the spectral shifts induced by exposure to oxygen, nitrogen or carbon monoxide.

Together, our results obtained with dried powders of lysozyme or haemoglobin demonstrate that carbohydrates enhance the stability of anhydrous proteins and permit the retention of physicochemical and biological activities after rehydration.

EXAMPLE 3

Our previous investigations of hemosomes as erythrocyte surrogates (Hayward et al., 1985) in

SUBSTITUTE SHEET

-16-

combination with our demonstration of protein stabilisation in the dry state by carbohydrate, prompted us to examine the capacity of sugars to stabilize artificial red blood cells in the dry state. We reasoned that the protective capacity of carbohydrate in this instance should be bimodal - prevention of the oxidation and denaturation of haemoglobin and the simultaneous retention of encapsulated haemoglobin within the internal aqueous compartment of the haemosome.

We found that inclusion of carbohydrates (both within the haemosome and in the external, bulk solution) prevented haem oxidation. Furthermore, rehydrated carbohydrate/haemosomes could be passed through a molecular sieve without diminution in the ratio of haemoglobin:phospholipid. All of the carbohydrates tested to-date have been effective in this capacity. Conversely, carbohydrate-free haemosomes became irreversibly aggregated and the phospholipid was readily separated from haemoglobin due to the former's relative buoyancy.

LITERATURE CITED

- Benesch, R., MacDuff, G. and Benesch, R.E/ (1965) Anal Biochem 11: 81 - 87.
- Bradbury, S.L. and Jakoby, W.B. (1972) Glycerol as an enzyme-stabilizing agent: effects on aldehyde dehydrogenase, Proc. Nat Acad Sci 69: 2373 - 2376.

SUBSTITUTE SHEET

-17-

Clegg, J.S. (1964) The control of emergence and metabolism by external osmotic pressure and the role of free glycerol in developing cysts of Artneia salina, J. Exp Biol 41: 879 - 892.

Crowe, J.H. and Clegg, J.S. (1973) "Anhydrobiosis:", Dowden, Hutchinson and Ross, Stroudsburg, P.A. p. 473.

Crowe, J.H, and Clegg, J.S. (1978) "Dry Biological Systems", Academic Press, NY, p. 357.

Crowe, J.H, Crowe, L.M. and Jackson S.A. (1983) Preservation of structural and functional activity in lyophilised sarcoplasmic reticulum, Arch Biochem Biophys. 220: 477 - 484.

Crowe, L.M. and Crowe, J.H. (1984) Effects of dehydration on membranes and membrane stabilisation at low water activities, in Chapman, D., ed., "Biological Membranes", Volume 5, Academic Press, London, pp. 57 - 102.

Crowe J.H., Crowe, L.M. and Chapman, D. (1984) Preservation of membranes in anhydrobiotic organisms: the role of trehalose, Science 223: 701 - 703.

Djordjevich, L. and Miller, I.F. (1980) Synthetic erythrocytes from lipid encapsulated hemoglobin, Exp. Hematol. 8: 584 -592.

Frigon, R.P. and Lee, J.C. (1972) The stabilization of calf-brain microtubule protein by sucrose, Arch. Biochem Biophys 153: 587 - 589.

Gaber, B., Yager, P., Sheridan, J.P. and Chang, E.L., (1983) Encapsulation of hemoglobin in phospholipid

SUBSTITUTE SHEET

-18-

vesicles, FEBS Lett. 153: 285 - 288.

Gerlsma, S.Y. (1968) J. Biol Chem 243: 957 - 963.

Hayward, J.A. Levine, D.M., Neufeld, L., Simon, S.R.,

Johnston, D.S. and Chapman D. (1985) Polymerized liposomes as stable oxygen-carriers, FEBS Lett 187: 261 - 266.

Madden, T.D., Bally, M.B., Hope, M.J., Cullis, P.R.,

Schieren, H.P. and Janoff, A.S. (1985) Protection of large unilamellar vesicles by trehalose during dehydration:

retention of vesicles contents, Biochem. Biophys Acta 817: 67 - 74.

Madin, K.A.C. and Crowe, J.H. (1975) Anhydrobiosis in nematodes: carbohydrate and lipid metabolism during dehydration. J. Exp. Zool. 193: 335 - 342.

Neucere, N.J. and St. Angelo, A.J. (1972) Physicochemical properties of peanut proteins in sucrose, Anal Biochem. 47: 80 - 89.

Szebeni, J., Di Iorio, E.E., Hauser, H. and Winterhalter, K.H. (1985) Encapsulation of hemoglobin in phospholipid vesicles: characterization and stability, Biochemistry 24: 2827 - 2832.

-19-

CLAIMS

1. A process for preserving a material having water-dependant structure comprising contacting the material with an aqueous solution of a polyhydroxy compound then removing water from the material.

2. A process according to claim 1 wherein the material is a protein, liposome or cell.

3. A process according to claim 1 wherein the material is an erythrocyte, oxyhaemoglobin, deoxyhaemoglobin or liposomes encapsulating oxyhaemoglobin or deoxyhaemaglonin.

4. A process according to claim 1 wherein the material encapsulates one or more spaces containing aqueous solutions or suspensions and wherein the process comprises contacting the material with, and introducing into one of more encapsulated spaces therein, an aqueous solution of a polyhydroxy compound then removing water from the material.

5. A process according to claim 1 wherein the polyhydroxy compound is a sugar or related polyol.

6. A process according to claim 5 wherein the polyhydroxy compound is selected from trioses, pentoses, hexoses, mono-, di- or oligosaccharides and related polyols.

-26-

7. A process according to claim 6 wherein the polyhydroxy compound is selected from glucose, galactose and non-reducing sugars.

8. A process according to claim 7 wherein the polyhydroxy compound is trehalose.

9. A dry formulation comprising a material having a water-dependant structure and a polyhydroxy compound in intimate admixture therewith.

10. A formulation according to claim 9 wherein the material is a protein, liposome or cell.

11. A formulation according to claim 9 wherein the material is an erythrocyte, oxyhaemoglobin, deoxyhaemoglobin or liposomes encapsulating oxyhaemoglobin or deoxyhaemoglobin.

12. A formulation according to claim 9 wherein the material encapsulates one or more spaces containing aqueous solutions or suspensions.

13. A formulation according to claim 9 wherein the polyhydroxy compound is a sugar or related polyols.

14. A formulation according to claim 13 wherein the polyhydroxy compound is selected from trioses, pentoses, hexoses, mono-, di- or oligosaccharides and related polyols.


- 21 -

15. A formulation according to claim 14 wherein the polyhydroxy compound selected from glucose, galactose and non-reducing sugars.

16. A formulation according to claim 15 wherein the polyhydroxy compound is trehalose.

1/1

Fig.1.



SUBSTITUTE SHEET



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁴ : C07K 3/00, 13/00, A61K 9/50 A61K 35/48 // A61K 37/04 C12N 9/36, 9/96	A3	(11) International Publication Number: WO 87/ 05300 (43) International Publication Date: 11 September 1987 (11.09.87)
(21) International Application Number: PCT/GB87/00143 (22) International Filing Date: 27 February 1987 (27.02.87) (31) Priority Application Number: 8604983 (32) Priority Date: 28 February 1986 (28.02.86) (33) Priority Country: GB (71) Applicant (for all designated States except US): BIOC-OMPATIBLES LIMITED [GB/GB]; 2 Gray's Inn Square, London WC1R 5AF (GB). (72) Inventors; and (75) Inventors/Applicants (for US only) : HAYWARD, James, Arthur [US/US]; 4 Chuck Court, Port Jefferson, NY 11777 (US). JOHNSTON, David, Samuel [GB/GB]; 65 Fitzjohns Avenue, Hampstead, London NW3 6PE (GB).		(74) Agents: CRESSWELL, Thomas, Anthony et al.; J.A. Kemp & Co., 14 South Square, Gray's Inn, London WC1R 5EU (GB). (81) Designated States: AT (European patent), BE (European patent), CH (European patent), DE (European patent), FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent), US. Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 22 October 1987 (22.10.87)
(54) Title: PROTEIN PRESERVATION (57) Abstract A process for preserving a material having water-dependant structure comprising contacting the material with an aqueous solution of a polyhydroxy compound then removing water from the material.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	ML	Mali
AU	Australia	GA	Gabon	MR	Mauritania
BB	Barbados	GB	United Kingdom	MW	Malawi
BE	Belgium	HU	Hungary	NL	Netherlands
BG	Bulgaria	IT	Italy	NO	Norway
BJ	Benin	JP	Japan	RO	Romania
BR	Brazil	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	LI	Liechtenstein	SN	Senegal
CH	Switzerland	LK	Sri Lanka	SU	Soviet Union
CM	Cameroon	LU	Luxembourg	TD	Chad
DE	Germany, Federal Republic of	MC	Monaco	TG	Togo
DK	Denmark	MG	Madagascar	US	United States of America
FI	Finland				

INTERNATIONAL SEARCH REPORT

PCT/GB 87/00143

International Application No

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ¹ According to International Patent Classification (IPC) or to both National Classification and IPC ⁴ C 07 K 3/00; C 07 K 13/00; A 61 K 9/50; A 61 K 35/18; IPC : //A 61 K 37/04; C 12 N 9/36; C 12 N 9/96																				
II. FIELDS SEARCHED <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black; margin: 5px 0;">Minimum Documentation Searched ¹</div> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 30%; border-bottom: 1px solid black;">Classification System</td> <td style="border-bottom: 1px solid black;">Classification Symbols</td> </tr> <tr> <td style="padding: 5px;">IPC ⁴</td> <td style="padding: 5px;">C 07 K 3/00; C 12 N 9/00; A 61 K 9/00</td> </tr> </table> <div style="text-align: center; border-top: 1px solid black; border-bottom: 1px solid black; margin: 5px 0;">Documentation Searched other than Minimum Documentation to the extent that such Documents are included in the Fields Searched ²</div>			Classification System	Classification Symbols	IPC ⁴	C 07 K 3/00; C 12 N 9/00; A 61 K 9/00														
Classification System	Classification Symbols																			
IPC ⁴	C 07 K 3/00; C 12 N 9/00; A 61 K 9/00																			
III. DOCUMENTS CONSIDERED TO BE RELEVANT ³ <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 10%; padding: 5px;">Category ⁴</th> <th style="width: 70%; padding: 5px;">Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²</th> <th style="width: 20%; padding: 5px;">Relevant to Claim No. ¹³</th> </tr> </thead> <tbody> <tr> <td style="text-align: center; vertical-align: top; padding: 5px;">X</td> <td style="padding: 5px;">US, A, 1855591 (L. WALLERSTEIN) 26 April 1932, see the whole document</td> <td style="text-align: center; vertical-align: top; padding: 5px;">1,2,5,6,7, 9,13,14</td> </tr> <tr> <td style="text-align: center; vertical-align: top; padding: 5px;">X</td> <td style="padding: 5px;">EP, A, 0111216 (TAKEDA CHEMICAL INDUSTRIES) 20 June 1984, see pages 59-60; example 11</td> <td style="text-align: center; vertical-align: top; padding: 5px;">1,2,5-10, 13,14</td> </tr> <tr> <td style="text-align: center; vertical-align: top; padding: 5px;">X</td> <td style="padding: 5px;">FR, A, 2532178 (ASAHI KASEI KOGYO K.K. & DAINIPPON PHARMACEUTICAL CO. LTD) 2 March 1984, see page 24, table 2; pages 28-32; claims 1-22</td> <td style="text-align: center; vertical-align: top; padding: 5px;">1,2,5-10, 13,14</td> </tr> <tr> <td style="text-align: center; vertical-align: top; padding: 5px;">X</td> <td style="padding: 5px;">Chemical Abstracts, volume 85, no. 3, 19 July 1976, (Columbus, Ohio, US), P. Labrude et al.: "Lyophilization of hemoglobin solutions. Study of protector compounds capable of preventing the formation of methemoglobin", see page 415, abstract 18011u, & J. Pharm. Belg. 1976, 31(2), 191-8</td> <td style="text-align: center; vertical-align: top; padding: 5px;">1,2,5-10, 13,14</td> </tr> <tr> <td style="text-align: center; vertical-align: top; padding: 5px;">X,Y</td> <td style="padding: 5px;">WO, A, 86/01103 (THE LIPOSOME COMPANY, INC.) 27 February 1986, see pages ./. .</td> <td style="text-align: center; vertical-align: top; padding: 5px;">1-14</td> </tr> </tbody> </table>			Category ⁴	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³	X	US, A, 1855591 (L. WALLERSTEIN) 26 April 1932, see the whole document	1,2,5,6,7, 9,13,14	X	EP, A, 0111216 (TAKEDA CHEMICAL INDUSTRIES) 20 June 1984, see pages 59-60; example 11	1,2,5-10, 13,14	X	FR, A, 2532178 (ASAHI KASEI KOGYO K.K. & DAINIPPON PHARMACEUTICAL CO. LTD) 2 March 1984, see page 24, table 2; pages 28-32; claims 1-22	1,2,5-10, 13,14	X	Chemical Abstracts, volume 85, no. 3, 19 July 1976, (Columbus, Ohio, US), P. Labrude et al.: "Lyophilization of hemoglobin solutions. Study of protector compounds capable of preventing the formation of methemoglobin", see page 415, abstract 18011u, & J. Pharm. Belg. 1976, 31(2), 191-8	1,2,5-10, 13,14	X,Y	WO, A, 86/01103 (THE LIPOSOME COMPANY, INC.) 27 February 1986, see pages ./. .	1-14
Category ⁴	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³																		
X	US, A, 1855591 (L. WALLERSTEIN) 26 April 1932, see the whole document	1,2,5,6,7, 9,13,14																		
X	EP, A, 0111216 (TAKEDA CHEMICAL INDUSTRIES) 20 June 1984, see pages 59-60; example 11	1,2,5-10, 13,14																		
X	FR, A, 2532178 (ASAHI KASEI KOGYO K.K. & DAINIPPON PHARMACEUTICAL CO. LTD) 2 March 1984, see page 24, table 2; pages 28-32; claims 1-22	1,2,5-10, 13,14																		
X	Chemical Abstracts, volume 85, no. 3, 19 July 1976, (Columbus, Ohio, US), P. Labrude et al.: "Lyophilization of hemoglobin solutions. Study of protector compounds capable of preventing the formation of methemoglobin", see page 415, abstract 18011u, & J. Pharm. Belg. 1976, 31(2), 191-8	1,2,5-10, 13,14																		
X,Y	WO, A, 86/01103 (THE LIPOSOME COMPANY, INC.) 27 February 1986, see pages ./. .	1-14																		
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>																				
IV. CERTIFICATION <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; border-bottom: 1px solid black; padding: 5px;">Date of the Actual Completion of the International Search</td> <td style="width: 50%; border-bottom: 1px solid black; padding: 5px;">Date of Mailing of this International Search Report</td> </tr> <tr> <td style="text-align: center; padding: 5px;">3rd September 1987</td> <td style="text-align: center; padding: 5px;">23 SEP 1987</td> </tr> <tr> <td style="border-bottom: 1px solid black; padding: 5px;">International Searching Authority</td> <td style="border-bottom: 1px solid black; padding: 5px;">Signature of Authorized Officer</td> </tr> <tr> <td style="text-align: center; padding: 5px;">EUROPEAN PATENT OFFICE</td> <td style="text-align: center; padding: 5px;">L. ROSSI</td> </tr> </table>			Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	3rd September 1987	23 SEP 1987	International Searching Authority	Signature of Authorized Officer	EUROPEAN PATENT OFFICE	L. ROSSI										
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report																			
3rd September 1987	23 SEP 1987																			
International Searching Authority	Signature of Authorized Officer																			
EUROPEAN PATENT OFFICE	L. ROSSI																			

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category*	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
	40-45; claims 1-61	
X,Y	Archives of Biochemistry and Biophysics", volume 242, no. 1, October 1985, Academic Press, Inc. L.M. Crowe et al.: "Preservation of freeze-dried liposomes by Trehalose", pages 240-247, see the whole document	1-14
X,Y	Chemical Abstracts, volume 103, no. 12, 23 September 1985, (Columbus, Ohio, US) T.D. Madden et al.: "Protection of large unilamellar vesicles by trehalose during dehydration: retention of vesicle contents", see page 316, abstract 92759a, & Biochim. Biophys. Acta 1985, 817(1), 67-74 cited in the application	1-14
Y	Biochemistry, volume 24, 1985, American Chemical Society, (Washington, US), J. Szebeni et al.: "Encapsulation of hemoglobin in phospholipid liposomes: Characterization and stability", pages 2827-2832, see pages 2827-2828 cited in the application	3,11
Y	Science, volume 223, no. 4637, 17 February 1984, American Association for the Advancement of Science, (Washington, US), J.H. Crowe et al.: "Preservation of membranes in anhydrobiotic organisms: the role of trehalose", pages 701-703, see the whole document cited in the application	1-14
Y	The Proceedings of Biotech '84 Europe, volume 1, American Chemical Society "World Biotech Report 1984", pages 379-390, see pages 382-385	1-3,5-8
P,X	WO, A, 86/03938 (THE REGENTS OF THE UNIVERSITY OF CALIFORNIA) 17 July 1986, see pages 18-21, claims 1-18	1-14
P,X	WO, A, 87/00196 (QUADRANT BIORESOURCES LTD.) 15 January 1987, see pages 34-35, claims 1-7	1-3,5-11, 13,14

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON

INTERNATIONAL APPLICATION NO. PCT/GB 87/00143 (SA 16358)

This Annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 17/09/87

The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A- 1855591		None	
EP-A- 0111216	20/06/84	EP-A, B 0037110	07/10/81
		JP-A- 56138248	28/10/81
		CA-A- 1162848	28/02/84
		US-A- 4517290	14/05/85
		JP-A- 57006363	13/01/82
		EP-A, B 0049898	21/04/82
		JP-A- 57118790	23/07/82
		CA-A- 1181342	22/01/85
		US-A- 4496658	29/01/85
FR-A- 2532178	02/03/84	JP-A- 59039829	05/03/84
		GB-A, B 2126588	28/03/84
		DE-A, C 3331003	01/03/84
		US-A- 4457916	03/07/84
		JP-A- 59059625	05/04/84
WO-A- 8601103	27/02/86	WO-A- 8601102	27/02/86
		EP-A- 0190315	13/08/86
		EP-A- 0191824	27/08/86
		JP-T- 62500101	16/01/87
		JP-T- 62500102	16/01/87
WO-A- 8603938	17/07/86	AU-A- 5318386	29/07/86
		EP-A- 0208764	21/01/87
		JP-T- 62501631	02/07/87
WO-A- 8700196	15/01/87	AU-A- 6136386	30/01/87
		EP-A- 0229810	29/07/87

For more details about this annex :
see Official Journal of the European Patent Office, No. 12/82